



Pulmonary Complication in Severe Malaria

Forman Erwin Siagian^{1*}

¹*Department of Parasitology and The Centre of Biomedic Research, Faculty of Medicine, Universitas Kristen Indonesia, Jakarta, Indonesia.*

Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/IJPR/2021/v8i130194

Editor(s):

(1) Prof. John Yahya I. Elshimali, Charles R. Drew University of Medicine and Science, USA.

Reviewers:

(1) Zaynab khorsheed Rasheed, Northern Technical University, Iraq.

(2) Rajapriya Manickam, St Joseph University, Nagaland, India.

Complete Peer review History: <https://www.sdiarticle4.com/review-history/73817>

Received 27 June 2021

Accepted 07 September 2021

Published 08 September 2021

Mini-review Article

ABSTRACT

Malaria is a potentially fatal disease caused by *Plasmodium* spp. Transmission occurs via the bite of female mosquito, *Anopheles* spp. Epidemiologically, global number of malaria patient are located in Southeast Asia and Africa. Until nowadays, millions of people still living in endemic area, with children and pregnant women are among the most vulnerable group in the population. Although there have been many advances in treatment and management, but the potential for harm remains; one of the example is lung involvement in patients with severe malaria. This paper aim to discuss briefly about lung derangement in the severe malaria and the inflammatory response related to the lung dysfunction. The severity of pulmonary impairment due to complications of malaria is determined not only by the initiation of antimalarial treatment but also by the hosts associated immune response

Keywords: *Blood protozoan; obligate intracellular; complication; ARDS; pulmonary vascular occlusion; inflammation.*

1. INTRODUCTION

The long known four species of blood protozoan parasites are *Plasmodium vivax*, *Plasmodium falciparum*, *Plasmodium malariae*, and

Plasmodium ovale, with the addition of the newer confirmed species, that can also infect humans, a zoonosis caused by *Plasmodium knowlesi* [1]. Other species of malaria also reported transmitted from animal to human (zoonosis), but

*Corresponding author: Email: forman.siagian@uki.ac.id;

the number reported is not much. Global malaria epidemiology study conducted by Qiao *et al.*, revealed its recent trend [2]. Though globally, the malaria age-standardized incidence rate (ASR) already decreased by an average 0.80% (95% confidence interval 0.58–1.02%) per year from 1990 to 2019; however, actually it slightly increased in 5 year period, from 3195.32 per 100 000 in 2015 to 3247.02 per 100 000 in 2019. The incidence rate of children under five years old was still higher than other age groups in the population. A total number of 40 countries had higher ASRs in the year 2019 compared to the data in the year 2015. The fastest augmentation in Cabo Verde (from only 2.02 per 100 000 and soaring very high to a number 597.00 per 100 000 population). After the year 2015, the ASRs in three regions: high-middle, middle and low-middle Socio-demographic Index regions began to arise and the trends remained in the up-rising pattern in the year 2019. Central, Western and Eastern Sub-Saharan Africa had the highest ASRs since 1990, and traveller number in Eastern and Western Sub-Saharan Africa increased by 31.24 and 7.58%, respectively, from 2017 to 2018. Especially, most countries with ASR over 10 000 per 100 000 had increase in traveller number from 2017 to 2018, with the highest change by 89.56% in Mozambique. In 2019, there were an estimated 229 million cases of malaria worldwide. The estimated number of malaria deaths stood at 409 000 in 2019. Children aged under 5 years are the most vulnerable group affected by malaria; in 2019, they accounted for 67% (274 000) of all malaria deaths worldwide [2]. Children and pregnant woman remain the most susceptible group in the population [3].

The classic triad of malaria is characterized by sign and symptom of high fever, anemia and splenomegaly. Without proper treatment, susceptible and or vulnerable people can immediately enter the danger phase, the severe malaria [4,5]. Severe malaria is a systemic and potentially lethal condition of illness which characterized by clearly marked dysfunction of one or more peripheral organs, such as the brain, the kidney and the lung. In the brain, it caused condition called cerebral malaria (CM), while in the kidney it caused acute kidney injury (AKI), and the last but not least, in the lung it caused acute respiratory distress syndrome (ARDS). This paper aims to review lung derangement in the context of the severe malaria and features of the inflammatory response are

related to the lung dysfunction observed in severe malaria.

2. EARLY PHASE OF MALARIA

This blood protozoa primarily transmitted via the proboscis of a female Anopheles mosquito that contain malarial sporozoite and then inserted them into susceptible humans [6,7]. Mosquitoes with a higher numbers of sporozoites in their salivary glands following blood-feeding activity, aided by its proboscis, are more possible to have created new infection in susceptible individuals; and in terms of transmission, of course it have done so faster compared to other mosquitoes with fewer number of parasites [3]. The mosquitoes with heavy parasite burdens are consequentially more likely to result in the hosts definite clinical blood stage infection.

After entering the bloodstream, at the beginning in the very early phase of infection, this blood protozoan then brought up to the liver and infect hepatocytes; this phase is called human liver stage [8]. Here, the parasite inside the liver cell change their forms several time; from sporozoites, it develop into schizonts which then burst, and release merozoites into the blood. This transition into a new form believe by many expert is a preparation of the parasite to infect its definite predilection, the erythrocyte and begin the massive reproduction, asexually. Sato *et al* mentioned that the essential role of early host interactions in the liver that may dampen the subsequent pro-inflammatory immune responses and influence the occurrence of vital organ failure such as the brain, and highlighting a novel checkpoint in the fatality of this pathology [9]. Even in this active development, how the parasites evade the hosts immune system while moving from the liver to the blood has still in the process of revealing. This obligate intracellular parasite actually dictate its will to a series of host cells as if rendered helpless and does not recognize the dangerous activities carried out by this *Plasmodium* spp. [10,11].

Following an infective mosquito bite, which is when the mosquito proboscis pierces the surface of the skin in search of food, sporozoites move with its gliding motility in the direction from the mosquito salivary gland cavities and then are discharged within the small vessels of the hosts skin [12]. After entering the blood stream, sporozoites carried away via the blood stream inside the blood vessels and enter the liver cells. Here they evolve into schizonts, which then burst

and release a huge number merozoites into the blood, leading to the clinical symptoms of malaria.[13,14].

Each invading sporozoite develop into a schizont that become very matures to yield up to 40 000 merozoites over a period of one to several weeks (long-term latent forms of 2 species of malaria belongs to *P. vivax* and also *P. ovale*; it called hypnozoites stage that may endure quiescent that clinically, it has been reported that it can last for months or years) [13,15]. In some species of parasite, particularly *P. vivax*, some sporozoites become hypnozoites. This form lies dormant in the liver and can reactivate leading to schizont formation and the ensuing symptoms of malaria in the absence of an infectious mosquito bite.16 According to Adekunle, based on their modeling of *P. vivax* primary infection and hypnozoite reactivation, that 90–96% of intra-erythrocyte infections emerge from the reactivation of dormant hypnozoite [16].

For most people, symptoms may occur in 10 days to 4 weeks after infection, although for a susceptible and immunological naive individual, he/she may feel ill as early as 1 week after the bite of mosquito or perhaps up to years later [5,17,18] Infected individual with clear clinical symptoms may suffer from cycles of malaria attacks. An attack usually starts with shivering and chills, followed by a high fever, followed by sweating and a return to normal temperature [19] Malaria signs and symptoms typically begin within a few weeks after being bitten by an infected mosquito. The species *P. falciparum* requires shorter duration to advance its erythrocytic stage compared to the other four species of human plasmodium while the longer ones are observed in *P. Malariae* [20,21].

Study conducted by Brasil *et al.*, showed us that on the hundreds of reported cases of imported *P. vivax* malaria patients hospitalized in the city of Rio de Janeiro, where there is no vector-borne transmission at all, the patients clinical data has made feasible for them to analyse the time-length of the incubation period between patients. The result made them realize that the time-period may be extended in certain cases. The heterogeneity in essential biological characteristics is traditionally appraised as one possible description to the origination of relapse in malaria and also to the differences in the time duration of the malaria incubation period among patients, a condition which can also be elucidated by the application of antimalaria chemoprophylaxis [22].

One of the most common condition found in malaria survivors is the repetitive occurrence of malaria attack [23] The term relapse is now used specifically to describe recurrences of malaria derived from persistent liver stages of the parasite (hypnozoites) [19] The number of sporozoites inserted and inoculated by the anopheline mosquito into the susceptible and vulnerable host is a crucial determinant of both the number and also the timing of relapses [24] The time intervals between each relapses reveal an exceptional periodicity with evidence is presented that the proportion of patients who unfortunately have repetitive relapses is relatively persistent. The internal factor which initiates dormant hypnozoites to wake up and become active and then escort to regular interval relapse in malaria is the condition of systemic febrile illness itself [25]. It is suggested that in an endemic areas of malaria where transmission of infection occurs heavily, then a large portion of the population bears dormant but latent hypnozoites which can be stimulated by an acute systemic infection, e.g., falciparum or vivax malaria. That is why, for returned travelers from malaria endemic area, they should consult to the doctor or healthcare providers of any travel in areas where definite malaria occurs during the past 12 months.

2.1 Uncomplicated and Complicated Malaria

Infection with malaria parasites may result in a wide variety of symptoms, ranging from absent or very mild symptoms to severe disease and even death, and the term uncomplicated and complicated is intended to make the division of symptoms a little easier, even though in daily practice, what was previously classified as uncomplicated malaria can very quickly turn into complicated malaria [18,26]. Delay in receiving treatment for uncomplicated malaria (UM) is often reported to increase the risk of developing severe malaria (SM) [27].

In patients with uncomplicated malaria, the classical subjective complain that infrequently observed during malaria episode may lasts 6–10 hours; the episodes consists of a cold phase (sensation of cold, shivering), and immediately followed by hot phase (fever, headaches, vomiting, and seizures in young children), and then ends in a sweating phase (sweats, return to normal temperature, tiredness) [17,28].

Traditionally based on the duration of asymptomatic period, classic malaria has a fever

pattern that appears over a period of time, which is the attacks occur on every second day with the type of tertian malaria parasites (*P. falciparum*, *P. vivax*, and *P. ovale*) and on every third day with the type of quartan parasite (*P. malariae*) [17,24,26]. More commonly, the patients present with a combination of the following symptoms: high fever, systemic malaise, chills, headaches, myalgia, nausea, and vomiting. During physical examination, doctor may found: the patients body temperatures rises, increased respiratory rate, massive perspiration, weakness of the muscle, light jaundice, enlargement of the liver and splenomegali (enlargement of the spleen). Splenomegali usually only occurs in people who live in endemic areas and a marker of chronicity, although it is possible that the patient actually does not show significant clinical symptoms [29].

3. ORIGIN OF SEVERITY OF THE LUNG DERANGEMENT

In malaria, the main finding of patients with falciparum malaria, the deadliest type of malaria infection, is the sequences of sequestration of infected erythrocytes within the microvasculature of the vital organs, e.g., the brain or the lung. Infected erythrocytes that contain falciparum malaria become more rigid due to the loss of deformability and flexibility properties [30-33]. Infected red blood cells actually underwent certain modifications right in its outer facet, a condition named knobs, and it seem to facilitate its adherence to the inner vascular endothelium and also perhaps to other erythrocytes, infected or non-infected ones [34-36]. The study conducted by Raj and Anderson revealed the role of erythrocyte (RBC) deformability and perfusate viscosity on lung segmental vascular resistance that in the end affect pulmonary hemodynamics [33] These findings suggest that decreases in erythrocyte deformability make it probably becomes stiffer intravascularly, so that it is easier to adhere to the vessel wall and then ends up with erythrocyte sequestration [34] Infected erythrocytes will never be the same as normal erythrocyte.

Adhesion actually supply better chance for the parasites ontogenesis in the condition that resembles microaerophilic venous atmosphere, cytoadherence contribute significantly in Plasmodiums immune-escape/evade mechanism, but at the same time also responsible to the pathogenesis of complication named severe malaria [37] Mature trophozoites and schizonts are sequestered and released

from the peripheral circulation due to adhesion of infected erythrocytes to the hosts endothelial cells. This chain of events directs the parasite to escape clearance by the spleen which actually recognizes the erythrocytes already loss its deformability [38]

Pathogenesis and worsening of the disease actually rely on adhesion to the inner part of microvasculatures endothelium, a condition termed *cytoadherence*, because adhesion contributed to occlusion of microvasculature, prevent sufficient blood supply to vital organs and leading to organ failure [39]. *Cytoadherence* can also describe adhesion of infected erythrocytes to uninfected erythrocytes, a phenomenon widely known as *rosetting* [37,40]. *Rosettes* were easily deranged by high shear forces inside the artery, but when it comes to the capillaries and post-capillary venules where the blood flow slows down dramatically, rosetting parasitized erythrocytes is easily bound to the inner microvasculatures endothelial cells and at the same time allows uninfected erythrocytes simultaneously to occlude vessels and reduce blood flow significantly [41] This condition is very danger to vital organs because permanent impediment of microvasculatures blood flow leading to the condition of reduced supply of oxygen and causing cells hypoxia, that lead to ischemia, and even metabolic disturbances. These serial of events surely caused permanent tissue damage, multi-organ failure and death in susceptible patient that suffer from severe malaria.

4. LUNG DERANGEMENT IN SEVERE MALARIA

Patients with severe falciparum, vivax, and knowlesi malaria may develop acute lung injury (ALI) and acute respiratory distress syndrome (ARDS) [42,43]. Ironically this conditon often developed just only several days after antimalarial medication was started, this condition could be due to the medication failure.⁴⁴ In south east Asia, decreased artesunate efficacy was already noted, and just as reported by Landre et al, combination of polymorphisms on the Kelch 13 (K13) gene and pharmacology of Artesunate itself was involved in therapeutic failure [44].

Epidemiologically, the rates ARDS incidence among severe falciparum malaria cases are 5% to 25% in adults and up to 29% in pregnant women, the condition of ARDS is scarce in the

group of young children.³² Condition of malignant respiratory distress evolves in up to 25% of adults and 40% of children. The underlying cause is a combination of several interrelated conditions including severe anemia, respiratory compensation of metabolic acidosis, noncardiogenic pulmonary edema and concomitant pneumonia [23,27,42,45].

Gas exchange is significantly impaired in patients with severe malaria.⁴⁶ Pulmonary edema is the most severe form of lung involvement among patient suffer from severe malaria, and this condition worsening the need for oxygen [47]. Increased alveolar capillary permeability leading to intravascular fluid loss into the lungs is the main pathophysiologic mechanism. This defines malaria as another cause of acute lung injury (ALI) and acute respiratory distress syndrome (ARDS).

In the case of severe falciparum malaria, it was associated with increased oxygen consumption but this was not related to disease severity [48]. Lactic acidosis did not result from inadequate overall macrocirculatory tissue oxygen delivery, but more likely from patchy microvascular perfusion abnormalities result in focal tissue hypoxia and combined with impaired hepatic clearance [43,45,46]. The haemodynamic status must kept optimized all the time to prevent the possibilities of decompensation in the period after antimalarial treatment [46]

Malaria related ARDS pathophysiology actually based on microvasculatures inflammatory-mediated responses that increased the capillars permeability and even to trigger dangerous endothelial damage that can happened diffusely; a condition that can persist even after post medication and after the clearance of the parasite with microvascular dysfunction likely contributing to impaired tissue perfusion [42,49,50].

The role of parasite sequestration in the human pulmonary microvasculature is indefinite due difficulties in establishing solid prove of sequestration, ethical issues and definitely the limitation of clinical report in human subjects [51-53]. It should not be forgotten also that clinico-epidemiologically, sequestration occurs intensely only in cases of *P. falciparum* infection, but less reported in the species *P. knowlesi*, and has not been shown credibly in *P. vivax* infection [51]. In Animal model, a group of scientist from Surabaya, Indonesia discovered all the

histopathological parameters of severity took place; including oedema, hemosiderin, thickened alveolar septa and inflammatory cell inside the lung from all infected mice with *P. berghei* that Exposed to Repeated Artemisinin [52]. The histological changes were severe. Those changes including diffuse oedema, increasing number of cellularity of the alveolar septa and thickening of the alveolar septa wall and massive infiltration of inflammatory cell in all affected areas of the lungs functional units with the existence of hemosiderin accumulation in septum inter-alveolar and degeneration of the bronchial epithelial. The finding of sequestered parasites and tissue damage in the lung infiltration clearly occurred in the lung [52].

In severe human malaria clinical case, ALI and ARDS be seen as part of a severe multi-system failure or may be the only main clinical attribute and often happens in condition of reduced parasitemia, soon after antimalarial medication given. In severe falciparum malaria, the roentgenographic presentations include diffuse interstitial edema, pulmonary edema, pleural effusion, and lobar consolidation.

Sometimes, doctors who treat patients with severe lung disorders due to malaria must make sure that fluid overload condition is prohibited and follow the old adage to keep them parched/dry because the earliest conclusive markers of ARDS are not yet available and do not forget that cases like this often occur in primary health care facilities with very limited equipment and resources [54,55]. Mechanical ventilation can save life in ALI/ARDS. Basic critical care facilities are increasingly available in tropical countries. The use of early lung-protective ventilation has helped to reduce mortality from malaria-induced ALI/ARDS, but unfortunately this action is not recommended for patient underwent permissive hypercapnia in unconscious severe malaria patient. This is due to the risk of elevated intracranial pressure and the potency of cerebral swelling that might occur in cerebral malaria. The best antimalarial treatment of severe malaria is IV artesunate. Even when proper treatment started immediately, and the parasite blood count showed improvement, the doctor must keep the patient closely observed in order to prevent the occurrence of post-treatment complication.

5. CONCLUSION

The best and optimal management of malaria-induced lung derangement includes early

recognition via a close observation to patients subjective complaint, clinical and laboratory findings and made the proper diagnosis, cito. Malaria cases must always be suspected if during anamnesis, the subject is complaining about acute febrile illness and just returned from a malaria-endemic region, e.g., a returning tourist or traveler or even just a short-term period visitor. Slide microscopy and/or the use of rapid antigen tests are standard diagnostic tools to make the correct diagnosis of malaria. Malaria must be treated with effective drugs, even current options are limited. Preventing clinical deterioration in malaria sufferers continues to be a challenge for health professionals.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Phillips M, Burrows J, Manyando C. Malaria. *Nat Rev Dis Primers*. 2017;3:17050. Available: <https://doi.org/10.1038/nrdp.2017.50>
2. Qiao L, Wenzhan J, Liangyu K, Jue L, Min L. Trends of the global, regional and national incidence of malaria in 204 countries from 1990 to 2019 and implications for malaria prevention, *Journal of Travel Medicine*, 2021;28(5):taab046. Available: <https://doi.org/10.1093/jtm/taab046>
3. Siagian FE. Malarial Related Myopathies: A Rhabdomyolysis Story. *Intl J Pathog Res*, 2020; 5(3): 39-51. Available: <https://doi.org/10.9734/ijpr/2020/v5i330137>
4. Souza M, Padua T, Henriques M. Multiple Organ Dysfunction During Severe Malaria: The Role of the Inflammatory Response; 2016. DOI: 10.5772/65348.
5. Guidelines for the Treatment of Malaria. 3rd edition. Geneva: World Health Organization; 2015. Available: <https://www.ncbi.nlm.nih.gov/books/NBK294440/>
6. Aleshnick M, Ganusov VV, Nasir G, Yenokyan G, Sinnis P. Experimental determination of the force of malaria infection reveals a non-linear relationship to mosquito sporozoite loads. *PLOS Pathogens*, 2020;16(5): e1008181. Available: <https://doi.org/10.1371/journal.ppat.1008181>
7. Churcher TS, Sinden RE, Edwards NJ, Poulton ID, Rampling TW, Brock PM, et al. Probability of Transmission of Malaria from Mosquito to Human Is Regulated by Mosquito Parasite Density in Naïve and Vaccinated Hosts. *PLoS Pathog*. 2017;13(1):e1006108. DOI: 10.1371/journal.ppat.1006108.
8. Mo AX, McGugan G. Understanding the Liver-Stage Biology of Malaria Parasites: Insights to Enable and Accelerate the Development of a Highly Efficacious Vaccine. *Am J Trop Med Hyg*. 2018;99(4):827-832. DOI: 10.4269/ajtmh.17-0895.
9. Sato Y, Ries S, Stenzel W, Fillatreau S, Matuschewski K. The Liver-Stage Plasmodium Infection Is a Critical Checkpoint for Development of Experimental Cerebral Malaria. *Front Immunol*. 2019;10:2554. DOI: 10.3389/fimmu.2019.02554.
10. Rogers KJ, Vijay R, Butler NS. Anti-malarial humoral immunity: the long and short of it. *Microbes Infect*. 2021;23(4-5):104807. DOI: 10.1016/j.micinf.2021.104807.
11. Yap, XZ, McCall, MBB, Sauerwein, RW. Fast and fierce versus slow and smooth: Heterogeneity in immune responses to Plasmodium in the controlled human malaria infection model. *Immunol Rev*. 2020;293:253–269. Available: <https://doi.org/10.1111/imr.12811>
12. Vaughan AM, Aly AS, Kappe SH. Malaria parasite pre-erythrocytic stage infection: gliding and hiding. *Cell Host Microbe*. 2008;4(3):209-18. DOI: 10.1016/j.chom.2008.08.010.
13. Maguire JD, Baird JK. Malaria in Aminoff MJ, Daroff RB (ed). *Encyclopedia of the Neurological Sciences* (2nd Edition). Academic Press, 2014;989-91. ISBN 9780123851581, Available: <https://doi.org/10.1016/B978-0-12-385157-4.00392-4>.
14. Beeson JG, Drew DR, Boyle MJ, Feng G, Fowkes FJ, Richards JS. Merozoite surface proteins in red blood cell invasion,

- immunity and vaccines against malaria. *FEMS Microbiol Rev.* 2016;40(3):343-72. DOI: 10.1093/femsre/fuw001.
15. Markus MB. Malaria: origin of the term hypnozoite. *J Hist Biol.* 2011;44(4):781-6. DOI: 10.1007/s10739-010-9239-3.
 16. Adekunle AI, Pinkevych M, McGready R, Luxemburger C, White LJ, et al. Modeling the Dynamics of Plasmodium vivax Infection and Hypnozoite Reactivation In Vivo. *PLOS Neglected Tropical Diseases*, 2015;9(3): e0003595. Available:https://doi.org/10.1371/journal.pntd.0003595
 17. Bria YP, Yeh CH, Bedingfield S. Significant symptoms and nonsymptom-related factors for malaria diagnosis in endemic regions of Indonesia. *Int J Infect Dis.* 2021;103:194-200. DOI: 10.1016/j.ijid.2020.11.177.
 18. Bartoloni A, Zammarchi L. Clinical aspects of uncomplicated and severe malaria. *Mediterr J Hematol Infect Dis.* 2012;4(1): e2012026. DOI: 10.4084/MJHID.2012.026.
 19. White NJ. Determinants of relapse periodicity in Plasmodium vivax malaria. *Malar J.* 2011;10:297. DOI: 10.1186/1475-2875-10-297.
 20. Orish V, Afutu L, Ayodele O, Likaj L, Marinkovic A, Sanyaolu A. A 4-Day Incubation Period of Plasmodium falciparum Infection in a Non-immune Patient in Ghana: A Case Report. *Open Forum Infect Dis.* 2019;6(1):ofy169. DOI: 10.1093/ofid/ofy169.
 21. Grande R, Antinori S, Meroni L. A case of Plasmodium malariae recurrence: recrudescence or reinfection?. *Malar J.* 2019;18:169. Available:https://doi.org/10.1186/s12936-019-2806-y
 22. Brasil P, de Pina Costa A, Pedro RS, da Silveira Bressan C, da Silva S, Tauil PL, Daniel-Ribeiro CT. Unexpectedly long incubation period of Plasmodium vivax malaria, in the absence of chemoprophylaxis, in patients diagnosed outside the transmission area in Brazil. *Malar J.* 2011;10:122. DOI: 10.1186/1475-2875-10-122.
 23. Kevin Baird J. Malaria caused by Plasmodium vivax: recurrent, difficult to treat, disabling, and threatening to life--the infectious bite preempts these hazards. *Pathog Glob Health.* 2013;107(8):475-9. DOI:10.1179/2047772413Z.000000000179
 24. White NJ, Imwong M. Relapse. *Adv Parasitol.* 2012;80:113-50. DOI: 10.1016/B978-0-12-397900-1.00002-5.
 25. de Araujo FCF, de Rezende AM, Fontes CJF, Carvalho LH, Alves de Brito CF. Multiple-Clone Activation of Hypnozoites Is the Leading Cause of Relapse in Plasmodium vivax Infection. *Plos One.* 2012;7(11): e49871. Available:https://doi.org/10.1371/journal.pone.0049871
 26. Milner DA Jr. Malaria Pathogenesis. *Cold Spring Harb Perspect Med.* 2018;8(1): a025569. DOI: 10.1101/cshperspect.a025569.
 27. Mousa A, Al-Taiar A, Anstey NM, Badaut C, Barber BE. The impact of delayed treatment of uncomplicated P. falciparum malaria on progression to severe malaria: A systematic review and a pooled multicentre individual-patient meta-analysis. *PLOS Medicine*, 2020;17(10): e1003359. Available:https://doi.org/10.1371/journal.pmed.1003359
 28. Laishram DD, Sutton PL, Nanda N. The complexities of malaria disease manifestations with a focus on asymptomatic malaria. *Malar J*, 2012;11:29. Available:https://doi.org/10.1186/1475-2875-11-29
 29. Kho S, Qotrunnada L, Leonardo L, Andries B, Wardani PAI, et al. Evaluation of splenic accumulation and colocalization of immature reticulocytes and Plasmodium vivax in asymptomatic malaria: A prospective human splenectomy study. *PLOS Medicine* 2021;18(5):e1003632. Available:https://doi.org/10.1371/journal.pmed.1003632
 30. Paul A, Ramdani G, Tatu U. Studying the rigidity of red blood cells induced by Plasmodium falciparum infection. *Sci Rep* 2019;9:6336 Available:https://doi.org/10.1038/s41598-019-42721-w
 31. Siagian FE, Isham D, Alfarabi M, Nainggolan S, Daroedono E, Novelyn S, et al. The profile of erythrocyte enlargement due to imported cases of Plasmodium vivax infection: its impact to the patient and the community. *Intl J Comm Med Publ Health*, 2020;7(7):2499-505. DOI:http://dx.doi.org/10.18203/2394-6040.ijcmph20202600

32. Taylor WRJ, Hanson J, Turner GDH, White NJ, Dondorp AM. Respiratory manifestations of malaria. *Chest*. 2012;142(2):492-505. DOI: 10.1378/chest.11-2655. PMID: 22871759.
33. Raj JU, Anderson J. Erythrocyte deformability and lung segmental vascular resistance: effect of hematocrit. *J Appl Physiol*. 1985;199;70(3):1386-92. DOI: 10.1152/jappl.1991.70.3.1386.
34. Huisjes R, Bogdanova A, van Solinge WW, Schiffelers RM, Kaestner L, van Wijk R. Squeezing for Life - Properties of Red Blood Cell Deformability. *Front Physiol*. 2018;9:656. DOI: 10.3389/fphys.2018.00656.
35. Kaul DK, Koshkaryev A, Artmann G, Barshtein G, Yedgar S. Additive effect of red blood cell rigidity and adherence to endothelial cells in inducing vascular resistance. *Am J Physiol Heart Circ Physiol*. 2008;295(4):H1788-93. DOI: 10.1152/ajpheart.253.2008. Epub 2008 Aug 29. PMID: 18757485.
36. Watermeyer JM, Hale VL, Hackett F, Clare DK, Cutts EE, Vakonakis I, et al. A spiral scaffold underlies cytoadherent knobs in Plasmodium falciparum-infected erythrocytes. *Blood*. 2016;127(3):343-51. DOI: 10.1182/blood-2015-10-674002.
37. Lee WC, Russell B, Rénia L. Sticking for a Cause: The Falciparum Malaria Parasites Cytoadherence Paradigm. *Front Immunol*. 2019;10:1444. DOI: 10.3389/fimmu.2019.01444.
38. Pivkin IV, Peng Z, Karniadakis GE, Buffet PA, Dao M, Suresh S. Biomechanics of red blood cells in human spleen and consequences for physiology and disease. *Proc Natl Acad Sci U S A*. 2016;113(28):7804-9. doi: 10.1073/pnas.1606751113.
39. Rajendran P, Rengarajan T, Thangavel J, Nishigaki Y, Sakthisekaran D, Sethi G, Nishigaki I. The vascular endothelium and human diseases. *Int J Biol Sci*. 2013;9(10):1057-69. DOI: 10.7150/ijbs.7502. PMID: 24250251; PMCID: PMC3831119.
40. Nunes-Alves, C. RIFINs promote rosette formation during malaria. *Nat Rev Microbiol*. 2015;13, 250. Available: <https://doi.org/10.1038/nrmicro3472>
41. Doumbo OK, Thera MA, Koné AK, Raza A, Tempest LJ, Lyke KE, Plowe CV, Rowe JA. High levels of Plasmodium falciparum rosetting in all clinical forms of severe malaria in African children. *Am J Trop Med Hyg*. 2009 Dec;81(6):987-93. doi: 10.4269/ajtmh.2009.09-0406.
42. Mohan A, Sharma SK, Bollineni S. Acute lung injury and acute respiratory distress syndrome in malaria. *J Vector Borne Dis*. 2008;45(3):179-93.
43. Pham TT, Verheijen M, Vandermosten L, Deroost K, Knoops S, Van den Eynde K, Boon L, Janse CJ, Opdenakker G, Van den Steen PE. Pathogenic CD8+ T Cells Cause Increased Levels of VEGF-A in Experimental Malaria-Associated Acute Respiratory Distress Syndrome, but Therapeutic VEGFR Inhibition Is Not Effective. *Front Cell Infect Microbiol*. 2017;7:416. DOI: 10.3389/fcimb.2017.00416.
44. Landre S, Bienvenu AL, Mialhes P, Abraham P, Simon M, Becker A, et al. Recrudescence of a high parasitaemia, severe Plasmodium falciparum malaria episode, treated by artesunate monotherapy. *Int J Infect Dis*. 2021;105:345-348. DOI: 10.1016/j.ijid.2021.02.080.
45. Taylor WR, Cañon V, White NJ. Pulmonary manifestations of malaria : recognition and management. *Treat Respir Med*. 2006;5(6):419-28. DOI: 10.2165/00151829-200605060-00007.
46. Anstey NM, Jacups SP, Cain T, Pearson T, Ziesing PJ, Fisher DA, Currie BJ, Marks PJ, Maguire GP. Pulmonary manifestations of uncomplicated falciparum and vivax malaria: cough, small airways obstruction, impaired gas transfer, and increased pulmonary phagocytic activity. *J Infect Dis*. 2002;185(9):1326-34. DOI: 10.1086/339885.
47. Moura G, Barcelos D, Epiphanyo S, Ortolan L. Physiopathology of Malaria-Associated Acute Respiratory Distress Syndrome. *Journal of Ancient Diseases & Preventive Remedies*. 2017;05(04). DOI: 10.4172/2329-8731.1000171.
49. Kingston HW, Ghose A, Rungpradubvong V. Does reduced oxygen delivery cause lactic acidosis in falciparum malaria? An observational study. *Malar J*. 2019;18: 97. Available: <https://doi.org/10.1186/s12936-019-2733-y>

50. White NJ. Malaria parasite clearance. *Malar J.* 2017;16(1):88. DOI: 10.1186/s12936-017-1731-1. Erratum in: *Malar J.* 2017 May 10;16(1):194. PMID: 28231817; PMCID: PMC5324257.
51. Barber BE, William T, Grigg MJ, Parameswaran U, Piera KA, et al. Parasite Biomass-Related Inflammation, Endothelial Activation, Microvascular Dysfunction and Disease Severity in Vivax Malaria. *PLOS Pathogens*, 2015;11(1):e1004558. Available: <https://doi.org/10.1371/journal.ppat.1004558>
52. Franke-Fayard B, Fonager J, Braks A, Khan SM, Janse CJ. Sequestration and tissue accumulation of human malaria parasites: can we learn anything from rodent models of malaria? *PLoS Pathog.* 2010;6(9):e1001032. DOI: 10.1371/journal.ppat.1001032.
53. Maslachah L, Widiyatno TV, Yustinasari LR. Sequestration and Histopathological Changes of the Lung, Kidney and Brain of Mice Infected with *Plasmodium berghei* that Exposed to Repeated Artemisinin. *Pakistan Veterinary Journal.* 2019;39(04): 499-504. ISSN 0253-8318, e-ISSN:2074-776
54. Jamrozik E, de la Fuente-Núñez V, Reis A. Ethical aspects of malaria control and research. *Malar J.* 2015;14:518. Available: <https://doi.org/10.1186/s12936-015-1042-3>
55. García-Laorden MI, Lorente JA, Flores C, Slutsky AS, Villar J. Biomarkers for the acute respiratory distress syndrome: how to make the diagnosis more precise. *Ann Transl Med.* 2017;5(14):283. DOI: 10.21037/atm.2017.06.49.
56. Elyazar IR, Hay SI, Baird JK. Malaria distribution, prevalence, drug resistance and control in Indonesia. *Adv Parasitol.* 2011;74:41-175. DOI:10.1016/B978-0-12-385897-900002-1.

© 2021 Siagian; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle4.com/review-history/73817>